

Supplementary Materials: A Stereocontrolled Protocol to Highly Functionalized Fluorinated Scaffolds through a Fluoride Opening of Oxiranes

Attila Márió Remete, Melinda Nonn, Santos Fustero, Ferenc Fülöp and Loránd Kiss





Figure 1. ¹H-NMR and ¹³C-NMR spectra of compound (±)-2.





Figure 2. ^1H -NMR and ^{13}C -NMR spectra of compound (\pm)-6.





Figure 3. ¹H-NMR and ¹³C-NMR spectra of compound (±)-10.





Figure 4. ^1H -NMR and ^{13}C -NMR spectra of compound (\pm)-13.





Figure 5. ¹H-NMR and ¹³C-NMR spectra of compound (±)-3.





Figure 6. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of compound (\pm)-4.






Figure 7. ¹H-NMR and ¹³C-NMR spectra of compound (±)-5.





Figure 8. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of compound (\pm)-7.






Figure 9. ¹H-NMR and ¹³C-NMR spectra of compound (±)-8.





Figure 10. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of compound (\pm)-11.








Figure 11. ¹H-NMR and ¹³C-NMR spectra of compound (±)-12.

Figure 12. ^1H -NMR and ^{13}C -NMR spectra of compound (\pm)-15.






Figure 13. ¹H-NMR and ¹³C-NMR spectra of compound (±)-14.





Figure 14. ¹H-NMR and ¹³C-NMR spectra of compound (±)-16.