

Design, Synthesis, and Biological Evaluation of 5,6,7,8-Tetrahydrobenzo[4,5]thieno[2,3-*d*]pyrimidines as Microtubule Targeting Agents

Farhana Islam ¹, Arpit Doshi ¹, Andrew J. Robles ^{2,3}, Tasdique M. Quadery ¹, Xin Zhang ¹, Xilin Zhou ¹, Ernest Hamel ⁴, Susan L. Mooberry ^{2,3,*} and Aleem Gangjee ^{1,*}

¹ Division of Medicinal Chemistry, Graduate School of Pharmaceutical Sciences, Duquesne University, 600 Forbes Avenue, Pittsburgh, PA 15282, USA; islamf@duq.edu (F.I.); doshia@duq.edu (A.D.); quaderyt@duq.edu (T.M.Q.); zhangx@duq.edu (X.Z.); zhoux@duq.edu (X.Z.)

² Department of Pharmacology, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229, USA; roblesa3@uthscsa.edu

³ Mays Cancer Center, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78229, USA

⁴ Molecular Pharmacology Branch, Developmental Therapeutics Program, Frederick National Laboratory for Cancer Research, Division of Cancer Treatment and Diagnosis, National Cancer Institute, National Institutes of Health, Frederick, MD 21702, USA; hamele@dc37a.nci.nih.gov

* Correspondence: mooberry@uthscsa.edu (S.L.M.); gangjee@duq.edu (A.G.)

Supplementary Materials

Docking scores were measured using Schrödinger Maestro suite (Schrödinger, LLC, New York, NY, 2020-2).

Table S1. Docking scores.

	Docking scores (kcal/mol)
1	-10.15
2	-10.11
3	-10.10
4	-10.89
5	-10.72
6	-10.52
7	-10.68
8	-10.55
9	-9.68
10	-10.42
11	-9.75
12	-10.40
13	-10.45
14	-9.75
Colchicine	-11.02




Figure S1. Superposition of 6BS2 pyrimidine ligand with compound **4**.