

Supporting Information

Simultaneous quantification of opioids in blood and urine by Gas chromatography-Mass spectrometer with modified dispersive solid-phase extraction technique

Sara Yasien¹, Ejaz Ali^{1*}, Mohsin Javed², Muhammad Muntazir Iqbal², Shahid Iqbal^{3*}, Hamad Alrbyawi⁴, Samar O. Aljazzar⁵, Eslam B. Elkaeed⁶, Ayed A. Dera⁷, Rami Adel Pashameah⁸, Eman Alzahrani⁹, Abd-ElAzim Farouk¹⁰

¹University College of Pharmacy, University of the Punjab, Lahore, Pakistan.

²Department of Chemistry, School of Science, University of Management and Technology, Lahore 54770, Pakistan.

³Department of Chemistry, School of Natural Sciences (SNS), National University of Science and Technology (NUST), H-12, Islamabad, 46000, Pakistan.

⁴Pharmaceutics and Pharmaceutical Technology Department, College of Pharmacy, Taibah University, Medina 42353, Saudi Arabia

⁵Department of Chemistry, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia.

⁶Department of Pharmaceutical Sciences, College of Pharmacy, AlMaarefa University, Riyadh 13713, Saudi Arabia.

⁷Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, King Khalid University, Abha, Saudi Arabia.

⁸Department of Chemistry, Faculty of Applied Science, Umm Al-Qura University, Makkah 24230, Saudi Arabia.

⁹Department of Chemistry, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia.

¹⁰Department of Biotechnology College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia.

***To whom corresponding should be addressed**

shahidgcs10@yahoo.com (Shahid Iqbal) and ejaz.pharmacy@pu.edu.pk (Ejaz Ali)



Figure. S1 *Dispersed phase in blood sample*



Figure. S2 *Dispersed phase in the urine sample*




Figure. S3 *Positive QC-Total ion chromatogram*




Figure. S4 *Negative QC-Total ion chromatogram*