

Supplementary Materials

Effect of Cationic Lipid Nanoparticle Loaded siRNA with Stearylamine against Chikungunya Virus

Manish Kumar Jeengar ^{1,2,†}, Mallesh Kurakula ^{3,4,†}, Poonam Patil ¹, Ashwini More ¹, Ramakrishna Sistla ^{3,*} and Deepti Parashar ^{1,*}

¹ ICMR-National Institute of Virology, 20-A, Dr. Ambedkar Road, Pune 411001, Maharashtra, India; manishkumarj@pharmacy.aims.amrita.edu (M.K.J.); poonamshewale07@rediffmail.com (P.P.); ashwini05.s@gmail.com (A.M.)

² Department of Pharmacology, Amrita School of Pharmacy, Amrita Vishwa Vidyapeetham, AIMS Health Sciences Campus, Kochi 682041, Kerala, India

³ CSIR-Indian Institute of Chemical Technology [CSIR-IICT], Hyderabad 500007, Telangana, India; mallesh_kurakula@yahoo.com

⁴ Product Development, CURE Pharmaceutical, Oxnard, CA 93033, USA.

* Correspondence: sistla@csiriict.in_(R.S.); deeptiparasharster@gmail.com (D.P.)

† These authors contributed equally to this work.




Figure S1. Agarose Gel electrophoresis results shows formation of Stable complexation of siRNA – F4 SLN; Lane 1 Naked siRNA; Lane 2–4 siRNA complexed with cationic SLN incubated at 4 °C; Lane 5 Blank SLN; Lane 6 DNA ladder; Lane 7–9, siRNA complexed with cationic SLN incubated at 37 °C.