

Cannabidiolic acid in hemp seed oil table spoon and beyond

Ersilia Nigro^{1,2,†}, Maria Tommasina Pecoraro^{1,†}, Marialuisa Formato¹, Sara Ragucci¹, Simona Piccolella¹, Marta Mallardo^{1,2}, Rosita Russo¹, Antimo Di Maro¹, Aurora Daniele^{2,3,*} and Severina Pacifico^{1,*}

¹ Dipartimento di Scienze e Tecnologie Ambientali, Biologiche, Farmaceutiche, Università della Campania "Luigi Vanvitelli", 81100, Caserta, Italy.

² CEINGE, Biotecnologie Avanzate Scarl, Napoli.

³ Dipartimento di Medicina Molecolare e Biotecnologie Mediche, "Federico II" Università degli Studi di Napoli, Napoli, 80131, Italy.

* Correspondence: severina.pacifico@unicampania.it; aurora.daniele@unina.it

† These authors equally contribute to the manuscript




Figure S1. TOF-MS and MS/MS spectra of CBDA detected in leaf hemp sample and cold-pressed hemp seed oil. Red arrows highlight the corresponding peaks in the TICs (Total Ion Chromatograms).




Figure S2. UHPLC-TOF-MS chromatogram of purified CBDA, acquired using the same experimental conditions of hemp leaf extract.




Figure S3. A) HPLC-UV chromatogram recorded at 266 ± 4 nm; B) HPLC-UV chromatogram recorded at 306 ± 4 nm; C) UV spectrum of purified CBDA.