

Non-Volatile Component and Antioxidant Activity: A Comparative Analysis between *Litsea Cubeba* Branches and Leaves

Wei Dai¹, Boyi Li², Yanli Xiong², Liping Dai¹, Yuan Tian³, Liangqian Zhang³, Qi Wang^{3,*} and Guoqiang Qian^{2,*}

¹ Teaching and Experimental Center, Guangdong Pharmaceutical University, Guangzhou 510006, China; dai_gdpu_2018@gdpu.edu.cn (W.D.)

² School of Chinese Medicine, Guangdong Pharmaceutical University, Guangzhou 510006, China

³ Key Laboratory of Xinjiang Phytomedicine Resource and Utilization, Ministry of Education, School of Pharmacy, Shihezi University, Shihezi 832003, China

* Correspondence: wq_ph@shzu.edu.cn (Q.W.); tgqqian@gdpu.edu.cn (G.Q.)

Figure S1. UPLC-Q-Orbitrap HRMS chromatograms of branches *Litsea cubeba* in negative ion mode

Figure S2. UPLC-Q-Orbitrap HRMS chromatograms of branches *Litsea cubeba* in positive ion mode

Figure S3. UPLC-Q-Orbitrap HRMS chromatograms of leaves *Litsea cubeba* in negative ion mode

Figure S4. UPLC-Q-Orbitrap HRMS chromatograms of leaves *Litsea cubeba* in positive ion mode




Figure S1. UPLC-Q-Orbitrap HRMS chromatograms of branches *Litsea cubeba* in negative ion mode




Figure S2. UPLC-Q-Orbitrap HRMS chromatograms of branches *Litsea cubeba* in positive ion mode

RT: 0.00 - 22.01




Figure S3. UPLC-Q-Orbitrap HRMS chromatograms of leaves *Litsea cubeba* in negative ion mode

RT: 0.00 - 22.01




Figure S4. UPLC-Q-Orbitrap HRMS chromatograms of leaves *Litsea cubeba* in positive ion mode