

The Importance of Substituent Position for Antibacterial Activity in the Group of Thiosemicarbazide Derivatives

Sara Janowska ^{1,*}, Joanna Stefańska ², Dmytro Khylyuk ³ and Monika Wujec ^{3,*}

¹ Department of Pathobiochemistry and Interdisciplinary Applications of Ion Chromatography, Biomedical Sciences, Medical University of Lublin, 1 Chodzki Street, 20-093 Lublin, Poland

² Department of Pharmaceutical Microbiology, Centre for Preclinical Research, Medical University of Warsaw, Banacha 1B Street, 02-097 Warsaw, Poland; joanna.stefanska@wum.edu.pl

³ Department of Organic Chemistry, Faculty of Pharmacy, Medical University, 4a Chodzki Street, 20-093 Lublin, Poland; dmytro.khylyuk@umlub.pl

* Correspondence: sara.janowska@umlub.pl (S.J.); monika.wujec@umlub.pl (M.W.)

Table of contents:

¹ H NMR spectra of thiosemicarbazide derivatives	2
¹³ C NMR spectra of thiosemicarbazide derivatives.....	5
IR spectra of thiosemicarbazide derivatives.....	7




Figure S1. The ^1H NMR of compound T4A.




Figure S2. The ^1H NMR of compound T5A.




Figure S3. The ^1H NMR of compound T8A.




Figure S4. The ^{13}C NMR of compound T4A.




Figure S5. The ^{13}C NMR of compound T5A.




Figure S6. The ^{13}C NMR of compound T8A.




Figure S7. The IR of compound T4A.




Figure S8. The IR of compound T5A.




Figure S9. The IR of compound T8A.