

New Low-Dimensional Organic–Inorganic Lead Halide Hybrid Systems Directed by Imidazo[1,5-*a*]pyridinium-Based Cation or Imines: Synthesis, Structures, Non-Covalent Interactions and Optical Properties

Olga Yu. Vassilyeva ^{1,*}, Elena A. Buvaylo ¹, Oksana V. Nesterova ², Alexandre N. Sobolev ³
and Dmytro S. Nesterov ^{2,*}

¹ Department of Chemistry, Taras Shevchenko National University of Kyiv,
64/13 Volodymyrska Str., 01601 Kyiv, Ukraine

² Centro de Química Estrutural, Instituto de Molecular Sciences, Instituto Superior Técnico,
Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

³ School of Molecular Sciences, M310, University of Western Australia, Perth, WA 6009, Australia

* Correspondence: vassilyeva@univ.kiev.ua (O.Y.V.); dmytro.nesterov@tecnico.ulisboa.pt (D.S.N.)

Supplementary Information




Figure S1. IR spectrum of [L1]_{2n}[Pb₂Cl₆]_{n∞}·nH₂O (1).




Figure S2. IR spectrum of $[PbBr_2(L2)]_{n\infty} \cdot 0.5nH_2O$ (2).




Figure S3. IR spectrum of $[PbBr_2(L3)]_2$ (3).




Figure S4. Selected fingerprint plots of the (N12, N13a) and (N22, N23a, major component) $[L1]^+$ cations in the crystal structure of **1** showing $X_i \cdots X_e$ interactions (where $X = \text{Cl}, \text{O}, \text{C}$ or H).




Figure S5. Selected fingerprint plots of the fragment $[\text{PbBr}_2(\text{L}2)]$ in the crystal structure of **2** showing $X_i \cdots X_e$ interactions (where $X = \text{Br}, \text{O}, \text{C}$ or H).




Figure S6. Selected fingerprint plots of the dinuclear molecule of **3** showing $X_i \cdots X_e$ interactions (where $X = \text{Br}, \text{N}, \text{C}$ or H).




Figure S7. Reduced density gradient surfaces (isovalue of 0.5) showing weak vdW interactions (green area) between aromatic fragments of the ligands L3 in the molecular structure of **3** (lead and bromine atoms were excluded from calculations).