

Table S2. The frequencies (%) and medians (\pm QD) of basic demographic and clinical characteristics of patients in relation to genotypes of the *PDGFA* gene polymorphisms.

Characteristics	rs1800814			rs2070958			rs62433334		
	AA	AG	GG	CC	CT	TT	CC	CG	GG
n	37	60	35	23	56	53	65	57	10
Male sex, n (%)	12 (32.43)	31 (51.67)	12 (34.29)	8 (34.78)	30 (53.57) ⁵	17 (32.08)	23 (35.38)	28 (49.12)	4 (40.00)
Age years, median \pm QD	46.50 \pm 4.00	45.00 \pm 6.25	46.50 \pm 5.50	45.00 \pm 5.50	45.50 \pm 6.00	47.00 \pm 5.50	46.00 \pm 6.50	46.50 \pm 3.75	43.00 \pm 5.00
BMI, median \pm QD	25.47 \pm 1.61	26.30 \pm 3.35	25.08 \pm 4.54	25.08 \pm 2.66	26.48 \pm 3.35	25.47 \pm 1.73	25.88 \pm 2.35	25.18 \pm 2.13	25.08 \pm 3.00
Cigarette smoking, n (%)	11 (29.73) ¹	6 (10.00)	5 (14.29)	4 (17.39)	6 (10.71)	12 (22.64)	9 (13.85)	13 (22.81)	0 (0.00)
Cigarettes n/day, median \pm QD	0.00 \pm 1.50	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00
Alcohol units/ week, median \pm QD	0.00 \pm 1.00	1.00 \pm 1.00	1.00 \pm 2.00	0.00 \pm 2.00	1.00 \pm 2.00	1.00 \pm 2.00	0.00 \pm 1.00	1.00 \pm 1.00	2.00 \pm 1.50
Hypertension, n (%)	7 (18.92)	3 (5.00)	8 (22.86)	3 (13.04)	8 (14.29)	7 (13.21)	12 (18.46)	5 (8.77)	1 (10.00)
Follow up therapy									
Physical therapy, n (%)	17 (45.95)	19 (31.67) ²	20 (57.14)	14 (60.87)	20 (35.71)	22 (41.51)	28 (43.08)	20 (35.09)	8 (80.00) ⁶
Manual therapy, n (%)	8 (21.62)	18 (30.00)	11 (31.43)	14 (60.87) ^{3, 4}	18 (35.71)	11 (20.75)	18 (27.69)	17 (29.82)	2 (20.00)
NSAIDs, n (%)	8 (21.62)	22 (36.67)	11 (31.43)	8 (34.78)	19 (33.93)	15 (28.30)	19 (29.23)	20 (35.09)	2 (20.00)

Legend: BMI, body mass index; NSAIDs, nonsteroidal anti-inflammatory drugs; QD, quartile deviation.

¹ AA vs AG, $P = 0.028$

² AG vs GG, $P = 0.015$

³ CC vs CT, $P = 0.019$

⁴ CC vs TT, $P = 0.000$

⁵ CT vs TT, $P = 0.024$

⁶ GG vs CC+CG, $P = 0.030$