

Curcumin-Rich Diet Mitigates Non-Alcoholic Fatty Liver Disease (NAFLD) by Attenuating Fat accumulation and Improving Insulin Sensitivity in Aged Female Mice Under Nutritional Stress

Gopal Lamichhane ¹, Da-Yeon Lee ¹, Rienna Franks¹, Femi Olawale ¹, Jong-Beom Jin ¹, Josephine M. Egan ², and Yoo Kim ^{1, *}

¹ Department of Nutritional Sciences, Oklahoma State University, Stillwater, OK 74078, USA

² Laboratory of Clinical Investigation, National Institute on Aging, Baltimore, MD 21224, USA

* Correspondence: yoo.kim@okstae.edu; Tel.: (optional; include country code; if there are multiple corresponding authors, add author initials)

Table S1. Sequence of primer used for real-time RT-PCR.

S.N.	Primer	Sequence (5'-3')
1.	ACC1	F-ATTGGGACCCCAGAGCTA
		R-CCCGCTCCTTCAACTTGCT
2.	ACC2	F-GGGCTCCCTGGATGACAAC
		R-TTCCGGGAGGAGTTCTGGA
3.	PPAR α	F-GCAGCTCGTACAGGTCATCA
		R-CTCTTCATCCCCAAGCGTAG
4.	PPAR γ	F-ATGCCAAAAATATCCCTGGTTTC
		GGAGGCCAGCATGGTGTAGA
5.	ChREBP	F-CTGGGGACCTAAACAGGAGC
		R-GAAGCCACCCTATAGCTCCC
6.	SREBP	F-GCGGTTGGCACAGAGCTT
		R-GGACTTGCTCCTGCCATCAG
7.	LPL	FTTTTCTGGGACTGAGGATGG
		GTCAGGCCAGCTGAAGTAGG
8.	β -actin	F-TACCACCATGTACCCAGGCA
		R-CTCAGGAGGAGCAATGATCCTGAT
9.	18S	F-CGATCCGAGGGCCTCATCA
		R-AGTCCCTGCCCTTTGTACACA

Figure S1. Detail of Western blot membrane.

