

A Randomized Controlled Prospective Cohort Study on the Efficacy of a Witch Hazel Extract Cream for the Eyelids and Eye Contour Area and a Cleansing Face Cream in Dermatitis of the Eyelids

Federica Veronese ¹, Elia Esposto ¹, Chiara Airoidi ², Nunzia Di Cristo ³, Pamela Paganini ³, Paola Savoia ^{3,*} and Elisa Zavattaro ³

Supplementary Material

Table S1. P-value obtained comparing mean values for T1 vs T0 and T2 vs T0, separately for treatment and controls using paired t-test.

Variable	Control	Treatment
	T1 vs T0, p-value	
DLQI	0.8853	0.2998
OSDI	0.6556	0.0495
NRS	0.2291	0.1173
EDSI total	0.3227	0.0731
	T2 vs T0, p-value	
DLQI	0.0940	0.0606
OSDI	0.1461	0.1093
NRS	0.0987	0.0100
EDSI total	0.1882	0.0092

Table S2. P-value obtained comparing median values for T1 vs T0 and T2 vs T0, separately for treatment and controls using Wilcoxon signed rank.

Variable	Control	Treatment
	T1 vs T0, p-value	
DLQI	>0.999	0.3062
OSDI	0.4443	0.0410
NRS	0.2813	0.1836
EDSI total	0.1436	0.0754
	T2 vs T0, p-value	
DLQI	0.1289	0.0699
OSDI	0.1445	0.0277
NRS	0.1318	0.0146
EDSI total	0.1201	0.0128