

First Direct Gravimetric Detection of PFAS Contaminants in Water. Combination with Electrical Measurements on the Same Device. Proof of Concepts

George R. Ivanov^{1,*}, Tony Venelinov², Yordan G. Marinov³, Georgi B. Hadjichristov³, Andreas Terfort⁴, Melinda David^{5,6}, Monica Florescu⁶ and Selcan Karakus⁷

¹ University of Architecture, Civil Engineering and Geodesy, University Lab "Nanoscience and Nanotechnology", 1 Hr. Smirnenki blvd., 1064 Sofia, Bulgaria; george@at-equipemnt.com

² University of Architecture, Civil Engineering and Geodesy; Laboratory for Water Analysis, 1 Hr. Smirnenki, blvd. , 1064 Sofia, Bulgaria; tvenelinov_fhe@uacg.bg

³ Institute of Solid State Physics, Bulgarian Academy of Sciences, 72, Tzarigradsko Chaussee blvd., Sofia, Bulgaria; ymarinov@issp.bas.bg and georgibh@issp.bas.bg

⁴ Department of Chemistry, Institute of Inorganic and Analytical Chemistry, Goethe University Frankfurt, Max-von-Laue-Str. 7, 60438 Frankfurt, Germany; aterfort@chemie.uni-frankfurt.de

⁵ Laboratory for Quality Control and Process Monitoring, University of Bucharest, 4-12 Elisabeta Blvd., 030018 Bucharest, Romania; melinda.david@s.unibuc.ro, melinda.david@unitbv.ro

⁶ Department of Fundamental, Prophylactic and Clinical Disciplines, Transilvania University of Brașov, Str. Universitatii no. 1, 500068 Brașov, Romania; florescum@unitbv.ro

⁷ Istanbul University-Cerrahpaşa, Faculty of Engineering, Department of Chemistry, (34320), İstanbul, Türkiye. selcan@iuc.edu.tr

* Correspondence: george@at-equipment.com Tel.: +359 885 465 535




Figure S1. Nitrogen adsorption-desorption for MIL-101(Cr) that was used in this study.




Figure S2. X-ray diffraction (XRD) for the powder raw MIL-101(Cr) material.