

Supplementary Materilas

Yield and composition variations of the milk from different camel breeds in Saudi Arabia

Amr A. El-Hanafy ^{1,2}, Yasser M. Saad ^{1,3}, Saleh A. Alkarim ¹, Hussein A. Almehdar ¹, Fuad M. Alzahrani ¹, Mo-hammed A. Almatry ¹, Vladimir N. Uversky ^{1,4,5,*}, and Elrashdy M. Redwan ^{1,6,*}

¹ Biological Sciences Department, Faculty of Sciences, King Abdulaziz University, P. O. Box 80203, Jeddah, 21589, Saudi Arabia

² Nucleic Acids Research Department, Genetic Engineering and Biotechnology Research Institute (GEBRI), City of Scientific Research and Technological Applications, Alexandria, Egypt, Borg El Arab, P. O. box. 21934.

³ Genetics Laboratory, National Institute of Oceanography and Fisheries, Egypt;

⁴ Institute for Biological Instrumentation of the Russian Academy of Sciences, Pushchino, Moscow region 142290, Russia

⁵ Department of Molecular Medicine and USF Health Byrd Alzheimer's Research Institute, Morsani College of Medicine, University of South Florida, Tampa, FL, USA

⁶ Therapeutic and Protective Proteins Laboratory, Protein Research Department, Genetic Engineering and Biotechnology Research Institute, City for Scientific Research and Technology Applications, New Borg EL-Arab 21934, Alexandria, Egypt

* Correspondence: lradowan@kau.edu.sa (E.M.R); vuversky@usf.edu (V.N.U.)



Figure S1: Ecotype of Saudi camel clans (Hamra, Safra, Majaheim and Wadha), milk of which was analyzed in this study.